

Monografia GSS 2008 – Rachide & Riabilitazione Multidisciplinare – Quarto Evidence Based Meeting

- 1 Delmas PD, Genant HK, Crans GG et al. Severity of prevalent vertebral fractures and the risk of subsequent vertebral and nonvertebral fractures: results from the MORE trial. *Bone*. 2003;33(4):522-32.
- 2 Diamond TH, Champion B, Clark WA. Management of acute osteoporotic vertebral fractures: a nonrandomized trial comparing percutaneous vertebroplasty with conservative therapy. *Am J Med*. 2003;114(4):257-65.
- 3 Diamond TH, Bryant C, Browne L, Clark WA. Clinical outcomes after acute osteoporotic vertebral fractures: a 2-year non-randomised trial comparing percutaneous vertebroplasty with conservative therapy. *Med J Aust*. 2006;184(3):113-7.
- 4 EPOS group. Incidence of vertebral fracture in europe: results from the European Prospective Osteoporosis Study (EPOS). *J Bone Miner Res*. 2002;17(4):716-24.
- 5 Fechtenbaum JC, Cropet S, Kolta S et al. The severity of vertebral fractures and health-related quality of life in osteoporotic postmenopausal women. *Osteoporos Int* 2005;16: 2175–2179.
- 6 Fink HA, Ensrud KE, Nelson DB et al. Disability after clinical fracture in post menopausal women with low bone density: the Fracture Intervention Trial. *Osteoporos Int* 2003;14:69–76.
- 7 Frost HM. Personal experience in managing acute compression fractures, their aftermath, and the bone pain syndrome, in osteoporosis. *Osteoporos Int*. 1998;8(1):13-5.
- 8 Genant HK, Wu CY, van Kuijk C, Nevitt MC. Vertebral fracture assessment using a semiquantitative technique. *J Bone Miner Res*. 1993;8(9):1137-48.
- 9 Grafe IA, Da Fonseca K, Hillmeier et al. Reduction of pain and fracture incidence after kyphoplasty: 1-year outcomes of a prospective controlled trial of patients with primary osteoporosis. *Osteoporosis Int*. 2005;16(12):2005-12.