

## Bibliografia

- 1 Roach J.W.: Adolescent idiopathic scoliosis. *Orthop Clin North Am* 1999, 30:353-365
- 2 Miller N.H.: Cause and natural history of adolescent idiopathic scoliosis. *Orthop Clin North Am* 1999, 30:343-352
- 3 Pehrsson K, Danielsson A, Nachemson A. Pulmonary function in adolescent idiopathic scoliosis: a 25 year follow up after surgery or start of brace treatment. *Thorax* 2001 May;56(5):388-393
- 4 Cotrel Y., Morel G.: La technique de l'EDF dans la correction des scolioses. *Rev Chir Orthop*, 1964; 50:59-75
- 5 Mauroy J.C., Stagnara P.: Resultats a long terme du traitement orthopedique lyonnais des scolioses inferieur a 50°. *Journées de la scoliose*, Lyon, 1979
- 6 Michel C.R., Caton J., Michel F.: Devenir a long terme des scolioses idiopathiques ayant subi un traitement orthopedique lyonnais. *Rev Chir Orthop* 1987, 73 (suppl 2) : 134-136
- 7 Gignac D, Aubin CE, Dansereau J, Poulin F, Labelle H. [A biomechanical study of new orthotic treatment approaches for the 3D correction of scoliosis]. *Ann Chir* 1998;52(8):795-800
- 8 Goldberg CJ, Moore DP, Fogarty EE, Dowling FE. Adolescent idiopathic scoliosis: the effect of brace treatment on the incidence of surgery. *Spine* 2001 Jan 1;26(1):42-47
- 9 Peterson LE, Nachemson AL. Prediction of progression of the curve in girls who have adolescent idiopathic scoliosis of moderate severity. Logistic regression analysis based on data from The Brace Study of the Scoliosis Research Society. *J Bone Joint Surg Am* 1995 Jun;77(6):823-827
- 10 Nachemson AL, Peterson LE. Effectiveness of treatment with a brace in girls who have adolescent idiopathic scoliosis. A prospective, controlled study based on data from the Brace Study of the Scoliosis Research Society. *J Bone Joint Surg Am* 1995 Jun;77(6):815-822
- 11 Rowe DE, Bernstein SM, Riddick MF, Adler F, Emans JB, Gardner-Bonneau D. A meta-analysis of the efficacy of non-operative treatments for idiopathic scoliosis. *J Bone Joint Surg Am* 1997 May;79(5):664-674
- 12 Danielsson AJ, Nachemson AL. Radiologic findings and curve progression 22 years after treatment for adolescent idiopathic scoliosis: comparison of brace and surgical treatment with matching control group of straight individuals. *Spine* 2001 Mar 1;26(5):516-525
- 13 Korovessis P, Kyrkos C, Piperos G, Soucacos PN. Effects of thoracolumbosacral orthosis on spinal deformities, trunk asymmetry, and frontal lower rib cage in adolescent idiopathic scoliosis. *Spine* 2000 Aug 15;25(16):2064-2071
- 14 Monticelli G., Illiano S., Corigliano A., Costanzo G.: comunicazione al Congresso ESDS-SRS, Amsterdam 1989.
- 15 Upadhyay SS, Nelson IW, Ho EK, Hsu LC, Leong JC. New prognostic factors to predict the final outcome of brace treatment in adolescent idiopathic scoliosis. *Spine* 1995 Mar 1;20(5):537-545
- 16 Pinto W.C., Avanzi O., Dezen E.: Common sense in the management of adolescent idiopathic scoliosis. *Orthop Clin North Am* 1994; 25:215-223
- 17 Ramirez N, Johnston CE 2nd, Browne RH, Vazquez S. Back pain during orthotic treatment of idiopathic scoliosis. *J Pediatr Orthop* 1999 Mar;19(2):198-201
- 18 Little DG, Song KM, Katz D, Herring JA. Relationship of peak height velocity to other maturity indicators in idiopathic scoliosis in girls. *J Bone Joint Surg Am* 2000 May;82(5):685-693
- 19 Aubin CE, Labelle H, Ruszkowski A, Petit Y, Gignac D, Joncas J, Dansereau J. Variability of strap tension in brace treatment for adolescent idiopathic scoliosis. *Spine* 1999 Feb 15;24(4):349-354
- 20 Wong MS, Mak AF, Luk KD, Evans JH, Brown B. Effectiveness and biomechanics of spinal orthoses in the treatment of adolescent idiopathic scoliosis (AIS). *Prosthet Orthot Int* 2000 Aug;24(2):148-162
- 21 Wong MS, Evans JH. Biomechanical evaluation of the Milwaukee brace. *Prosthet Orthot Int* 1998 Apr;22(1):54-67
- 22 Caton J., Michel C.R., Fiquet A., Jouvinroux P. : Le traitement orthopedique des scolioses lombaires et thoracolumbaires par l'orthose rachidienne à 3 valves. *Rev Chir Orthop* 1987, 73 (suppl 2) :138-142
- 23 Michel C.R., Caton J., Allègre G.: Place de l'orthèse rachidienne a 4 valves dans le traitement orthopedique des scolioses. *Rev Chir Orthop* 1983, 69 (suppl 2) : 135-140
- 24 Carr W.A., Moe J.H., Winter R.B., Lonstein J.E.: Treatment of idiopathic scoliosis in the Milwaukee brace, Long-term results. *J Bone Joint Surg*, 1980, 62-A:599-612
- 25 Edmonson A., Smith G.R.: Long-term follow-up study of Milwaukee brace treatment in patients with idiopathic scoliosis. *Spine*, 1983, 7:10-11.
- 26 Mellencamp D.D., Blount W.P., Anderson A.J.: Milwaukee brace treatment of idiopathic scoliosis: late results. *Clin Orthop* 1977, 126:47-57
- 27 Salanova C.: Late results of Milwaukee brace treatment of idiopathic scoliosis. *Orthop Trans* 1986, 10:2-5
- 28 Schuffelbarger H.L., Keiser R.P., King W.: Nonoperative treatment of idiopathic scoliosis: a 10 years study. *Orthop Trans* 1983, 7:11-13
- 29 Emans J.B., Kaelin A., Bancel P., Hall J.E., Miller M.E.: The Boston bracing system for idiopathic scoliosis. Follow-up results in 295 patients. *Spine* 1996, 11:792-801

- 30 Wiley JW, Thomson JD, Mitchell TM, Smith BG, Banta JV.  
Effectiveness of the Boston brace in treatment of large curves in adolescent idiopathic scoliosis. *Spine* 2000 Sep 15;25(18):2326-2332
- 31 Labelle H, Dansereau J, Bellefleur C, Poitras B. Three-dimensional effect of the Boston brace on the thoracic spine and rib cage.  
*Spine* 1996 Jan 1;21(1):59-64
- 32 Cheneau J.: L'orthèse de Munster. *Ann Med Phys*, 1981, 24 :377-391
- 33 Matussek J, Mellerowicz H, Klockner C, Sauerlandt B, Nahr K, Neff G.  
[2- and 3-dimensional correction of scoliosis by corset treatment. Optimized conservative therapy of idiopathic scoliosis with the improved Cheneau corset]. *Orthopade* 2000 Jun;29(6):490-499
- 34 Perie D, Sales De Gauzy J, Sevely A, Hobatho MC. In vivo geometrical evaluation of Cheneau-Toulouse-Munster brace effect on scoliotic spine using MRI method. *Clin Biomech (Bristol, Avon)* 2001 Feb;16(2):129-137
- 35 Howard A, Wright JG, Hedden D. A comparative study of TLSO, Charleston, and Milwaukee braces for idiopathic scoliosis.  
*Spine* 1998 Nov 15;23(22):2404-2411
- 36 Hopf C, Heine J. [Long-term results of the conservative treatment of scoliosis using the Cheneau brace]. *Z Orthop Ihre Grenzgeb* 1985 May;123(3):312-322
- 37 von Deimling U, Wagner UA, Schmitt O. [Long-term effect of brace treatment on spinal decompensation in idiopathic scoliosis. A comparison of Milwaukee brace--Cheneau corset]. *Z Orthop Ihre Grenzgeb* 1995 May;133(3):270-273
- 38 Katz DE, Richards BS, Browne RH, Herring JA. A comparison between the Boston brace and the Charleston bending brace in adolescent idiopathic scoliosis. *Spine* 1997 Jun 15;22(12):1302-1312
- 39 Noonan KJ, Dolan LA, Jacobson WC, Weinstein SL. Long-term psychosocial characteristics of patients treated for idiopathic scoliosis. *J Pediatr Orthop* 1997 Nov;17(6):712-717
- 40 Climent JM, Sanchez J. Impact of the type of brace on the quality of life of Adolescents with Spine Deformities. *Spine* 1999 Sep 15;24(18):1903-1908
- 41 Di Raimondo C.V., Green N.E.: Brace-wear compliance in patients with adolescent idiopathic scoliosis. *J Pediatr Orthop* 1988, 8:143-146
- 42 Mehta M.H.: The rib vertebral angle in the early diagnosis between resolving and progressive infantile scoliosis. *J Bone Joint Surg* 1972, 54-B:230-243